

# ON NON-NEGATIVELY CURVED METRICS ON OPEN FIVE-DIMENSIONAL MANIFOLDS

VALERY MARENICH AND MIKAEL BENGTSSON

**ABSTRACT.** Let  $V^n$  be an open manifold of non-negative sectional curvature with a soul  $\Sigma$  of co-dimension two. The universal cover  $\tilde{N}$  of the unit normal bundle  $N$  of the soul in such a manifold is isometric to the direct product  $M^{n-2} \times R$ . In the study of the metric structure of  $V^n$  an important role plays the vector field  $X$  which belongs to the projection of the vertical planes distribution of the Riemannian submersion  $\pi : V \rightarrow \Sigma$  on the factor  $M$  in this metric splitting  $\tilde{N} = M \times R$ . The case  $n = 4$  was considered in [GT] where the authors prove that  $X$  is a Killing vector field while the manifold  $V^4$  is isometric to the quotient of  $M^2 \times (R^2, g_F) \times R$  by the flow along the corresponding Killing field. Following an approach of [GT] we consider the next case  $n = 5$  and obtain the same result under the assumption that the set of zeros of  $X$  is not empty. Under this assumption we prove that both  $M^3$  and  $\Sigma^3$  admit an open-book decomposition with a bending which is a closed geodesic and pages which are totally geodesic two-spheres, the vector field  $X$  is Killing, while the whole manifold  $V^5$  is isometric to the quotient of  $M^3 \times (R^2, g_F) \times R$  by the flow along corresponding Killing field.

## 1. Introduction

Let  $(V^n, g)$  be a complete open Riemannian manifold of non-negative sectional curvature. Remind that as follows from [CG] and [P] an arbitrary complete open manifold  $V^n$  of non-negative sectional curvature contains a closed absolutely convex and totally geodesic submanifold  $\Sigma$  (called a soul) such that the projection  $\pi : V \rightarrow \Sigma$  of  $V$  onto  $\Sigma$  along geodesics normal to  $\Sigma$  is well-defined and is a Riemannian submersion (see also [CaS]). The (vertical) fibers  $F_P = \pi^{-1}(P), P \in \Sigma$  of  $\pi$  define a metric foliation in  $V$  and two distributions: a vertical  $\mathcal{V}$  distribution of subspaces tangent to fibers and a horizontal distribution  $\mathcal{H}$  of subspaces normal to  $\mathcal{V}$ . For an arbitrary point  $P$  on  $\Sigma$ , an arbitrary geodesic  $\gamma(t)$  on  $\Sigma$  and arbitrary vector field  $V(t)$  which is parallel along  $\gamma$  and normal to  $\Sigma$  the following

$$(1) \quad \Pi(t, s) = \exp_{\gamma(t)} s V(t)$$

are totally geodesic surfaces in  $V^n$  of zero curvature, i.e., flats.

When  $\dim \Sigma = 1$  or  $\text{codim}(\Sigma) = 1$  the manifold  $V^n$  is locally isometric to the direct product of  $\Sigma$  and Euclidean space of a complementary dimension and of non-negative curvature. Study of the next case  $\text{codim}(\Sigma) = 2$  was began in [M1], where we noted that the manifold  $V^4$  or is a direct product when the holonomy of the normal bundle of  $\Sigma$  in  $V$  is trivial, or the holonomy group acts transitively on normal vectors, every geodesic normal to  $\Sigma$  is a ray and (1) holds. The metric structure in this case might be more complicated. In [GT] the authors consider four-dimensional manifolds diffeomorphic to direct products  $M^2 \times R^2$  and prove the following.

---

1991 *Mathematics Subject Classification.* 53C20, 53C21. Supported by the Faculty of Natural Sciences of the Hogskolan i Kalmar, (Sweden).

*Key words and phrases.* open manifolds, non-negative curvature.

**Theorem A, [GT].** *Every non-negatively curved metric on  $M^2 \times R^2$  is isometric to a Riemannian quotient of the form  $((M^2, g_0) \times (R^2, g_F) \times R)/R$ . Here  $R$  acts diagonally on the product by the flow along Killing vector fields on  $(M^2, g_0)$  and  $(R^2, g_F)$  and by translations on  $R$ .*

The very important role in the proof of the Theorem A plays the vector field  $X$  which is the projection of the vertical vector field in the universal cover  $\tilde{N}$  of the boundary of some metric  $s$ -tube  $N$  of the soul on the "horizontal" factor  $M$  in the metric splitting  $\tilde{N} = M \times R$ , see below. In the four-dimensional case this vector field  $X$  restricted to  $M^2$  always has zeros since  $M^2$  is a two-dimensional sphere. In our case the soul  $\Sigma$  of  $V$  and  $M^3$  are three-dimensional spheres, and hence,  $X$  might be nowhere zero as the following simple example shows. Let  $h : S^3 \rightarrow S^2$  be the Hopf bundle, i.e., the factoring of a unit sphere  $S^3$  in the complex plane  $C^2$  by  $S^1$  action - multiplication by complex numbers of absolute value 1. Consider  $V^5$  which is the the quotient of the direct product of  $S^3 \times R^2 \times S^1$ , where the  $S^1$  acts on  $R^2 = C$  by rotations, i.e., again, multiplication in  $C^1$  by unit complex numbers. Then for the manifold  $V^5 = S^3 \times R^2 \times S^1/S^1$  the vector field  $X$  is nowhere zero.<sup>1</sup> The objective of this note is to expand an approach from [GT] to the case of non-negatively curved five-dimensional  $V^5$  diffeomorphic to a direct product  $S^3 \times R^2$  under the following assumption.

**Assumption 1.** *The set of zeros of the vector field  $X$  is not empty.*

Our main result is very similar to the Theorem A above.

**Theorem B.** *Let  $V^5$  be an open manifold of non-negative sectional curvature and diffeomorphic to  $S^3 \times R^2$ . Assume that the vector field  $X$  has non-empty zero set  $\mathcal{Z}$ . Then  $\mathcal{Z}$  is a closed geodesic and the manifold  $M^3$  admits a singular foliation - "open-book decompositions" by totally geodesic and isometric to each other horizontal two-dimensional spheres  $S^2(\psi)$ , where the singular set of this decompositions - "bindings", equal the closed geodesic  $\mathcal{Z}$ . The flow along Killing field  $X$  acts as "turning pages" in this open-book decomposition, while  $V^5$  itself is isometric to a Riemannian quotient of the form  $M^3 \times (R^2, g_F) \times R/R$  with  $R$  acting diagonally on the product by the flow along Killing vector fields on  $M^3$  and  $(R^2, g_F)$  and by translations on  $R$ . The Riemannian submersion  $\pi : V^5 \rightarrow \Sigma$  conveys the open-book decomposition of  $M^3$  to a similar open-book decomposition of  $\Sigma$  with the pages  $S^2(\psi)$  isometric to  $S^2(\psi)$ .*

In the same way as Theorem A in [GT] our Theorem B follows from the fact that the vector field  $X$  on  $N$  is Killing for every  $s$ , where  $N$  is the boundary of  $s$ -metric neighborhood of the soul  $\Sigma$  in  $V$ , see Theorem 3 below. Thus, after proving Theorem 3, we complete the proof of Theorem B by referring to the corresponding arguments from [GT], see section 5.

Note that the general case of five-dimensional open manifold  $V^5$  with a soul of codimension 2 can be reduced to the one under consideration as follows. First, we note that if the fundamental group of  $\Sigma$  (which is isomorphic to that of  $V$ ) is not finite, the universal cover  $\tilde{V}$  contains a straight line in the universal cover  $\tilde{\Sigma}$  of the soul. Then both  $\tilde{V}$  and  $\tilde{\Sigma}$  split into direct products, and the case is reduced to the already studied one of open four-dimensional manifolds. When the fundamental group of  $\Sigma$  is finite the universal cover  $\tilde{\Sigma}$  is diffeomorphic to a sphere  $S^3$  due to the non-negativity of the curvature. Next: because an arbitrary vector bundle over simply connected  $S^3$  is, obviously, trivial we see that an investigation of the metric structure of an arbitrary  $V^5$  with a soul of codimension 2 is reduced to the case when  $V^5$  is diffeomorphic to the direct product  $S^3 \times R^2$ .<sup>2</sup>

Below we assume that the holonomy of the normal bundle is not trivial, for otherwise by a direct product theorem from [M1,4] the manifold  $V$  is a metric product.

<sup>1</sup>For corresponding  $M^3$  and  $S^3$  the one-form given by the scalar product with  $X$  is a (nowhere degenerated) contact form  $\alpha$  with  $\alpha \wedge d\alpha$  - the volume form.

<sup>2</sup>The case when five-dimensional  $V^5$  has a soul of codimension 3 we considered in [M5].

## 2. Vector field $X$ and its zeros

Fix some positive  $s_0$  smaller than a focal radius of  $\Sigma$  in  $V$ . For some  $s < s_0$  denote by  $N\Sigma(s)$ , or simply by  $N$ , the boundary of an  $s$ -neighborhood of  $\Sigma$ . Due to our choice it is a smooth manifold. It consists of all points  $Q(P, V) = \exp_P(sV)$ , where  $P$  is a point on  $\Sigma$  and  $V$  is a unit vector normal to  $\Sigma$  at  $P$ .

**Lemma 1.**  *$N(s)$  has non-negative curvature if  $s$  is sufficiently small.*

*Proof.* This follows from the Gauss equations and the fact that  $N(s)$  bounds a convex subset in a manifold  $V$  of non-negative curvature. The last is obviously true when the holonomy of the (trivial) normal bundle  $\nu\Sigma$  of the soul is trivial, i.e., all parallel translations along closed curves in  $\Sigma$  acts identically on vectors normal to  $\Sigma$  because then  $V$  is isometric to the direct metric product  $\Sigma \times (R^2, h)$ , see [M1]. If the holonomy is not trivial, then all normal vectors are so called ray directions, and  $N(s)$  coincides with the boundary  $\partial C_s$  of an absolutely convex set constructed in [CG], see again [M1]. The Lemma 1 is proved.

**Lemma 2.** *The universal cover  $\tilde{N}$  of  $N(s)$  is isometric to the direct product  $(M, g) \times R$ , where  $M$  is diffeomorphic to  $S^3$ . The composition of a covering map and a submersion  $\pi$  provides a diffeomorphism between an arbitrary factor  $M$  and the soul  $\Sigma$  which we denote by  $\pi^M : M \rightarrow \Sigma$ .<sup>3</sup>*

*Proof.* This follows from the fact that  $N(s)$  is diffeomorphic to the trivial circle bundle over three-dimensional sphere  $\Sigma$ , i.e., has an infinite cycle fundamental group generated by a homotopy class of a fiber. Then by standard arguments the universal cover  $\tilde{N}(s)$  admits a straight line, and hence by Toponogov splitting theorem is isometric to the direct product  $(M, g) \times R$ .

Denote by  $E$  the unit vector field in  $N(s)$  tangent to the projections of straight lines (i.e.,  $R$ -factor) from  $\tilde{N}$  to  $N$ . By  $W$  we denote the (vertical) vector field on  $N$  which is the speed of the natural  $S^1$ -action on  $N$  given by rotations in a positive direction of a normal vectors to  $\Sigma$  as follows: for  $Q = Q(P, V)$  denote by  $Q_\phi = Q_\phi(P, V) = Q(P, V_\phi)$ , where  $V_\phi$  is  $V$  rotated by the angle  $\phi$  in the bundle of unit normals to  $\Sigma$  in  $V$  (which is correctly defined since the bundle is topologically trivial). Finely, denote by  $X$  the vector field on  $N$  which is the component of  $W$  normal to  $E$ .<sup>4</sup>

$$(2) \quad X = W - (W, E)E.$$

Note, that  $N$  naturally inherits from  $V$  a horizontal distribution  $\mathcal{H}$ , while the vector field  $W$  belongs to the vertical distribution. If by  $M$  we denote an image of some  $(M, g)$ -factor in the direct metric product  $\tilde{N}$  under the projection  $pr : \tilde{N} \rightarrow N$ , then (the restriction of)  $E$  on  $M$  would be the unit vector field of normals to  $M$ ,  $X$  is a vector field tangent to  $M$ , while another vector field  $Y$  tangent to  $M$  would be a horizontal if and only if it is normal to  $X$ . In particular, the tangent subspace  $T_Q M$  is horizontal  $\mathcal{H}_Q$  if and only if  $X(Q) = 0$ .

Note that the vector field  $W$  in  $N$  is never tangent to any of the  $M$ -factor, or (equivalently) never orthogonal to  $E$ . Indeed, if so then some homotopically non-trivial closed geodesic  $\Gamma(s)$  in  $N$  which is the images of a straight line in the universal cover  $\tilde{N} = M \times R$ , would be horizontal at some point, and therefore, horizontal everywhere, which obviously can not be homotopically non-trivial in  $N$ . To see this denote by  $\bar{\Gamma}(s)$  its image under  $\pi$  in  $\Sigma$ , and by  $V(s)$  the normal vector field of vertical geodesics connecting  $\bar{\Gamma}(s)$  and  $\Gamma(s)$ . Since  $\Sigma$  is simply connected there exists a disk  $D$  in  $\Sigma$  with a boundary  $\bar{\Gamma}$  and extension of the vector field  $V$  over  $D$  (because the restriction of a normal bundle to  $D$  is trivial). The vertical lift of  $D$  along this extension will provide us a disk in  $N$  with a boundary  $\Gamma$  implying that  $\Gamma$  is contractible in  $N$ . The obtained contradiction proves that  $E$  is never horizontal, or that the map  $\pi^M : M \rightarrow \Sigma$  from any of the image  $M$  of a factor in the direct product  $\tilde{N} = (M^3, g) \times R$  into the soul  $\Sigma$  is a diffeomorphism. The Lemma 2 is proved.<sup>5</sup>

<sup>3</sup>Note, that this statement and forthcoming (3) both are true for an arbitrary  $V^n$  with simply connected soul of codimension two.

<sup>4</sup>Note, that our  $X$  is different from similar  $X$  of [GT].

<sup>5</sup>This also fills the gap in the arguments from Lemma 2.1 in [GT].

By definition the differential of the diffeomorphism  $\pi^M : M \rightarrow \Sigma$  is an isometry on the subspace of horizontal vectors, i.e., on the subspace in  $T_Q M$  normal to  $X$  (or on the whole  $T_Q M$  if  $X(Q) = 0$ ), while

$$(3) \quad \|d\pi_Q^M(X)\| = \cos(\alpha(Q))\|X\|,$$

where  $\alpha(Q)$  denotes the angle at the point  $Q \in M$  between vectors  $E$  and  $W$ . The map  $\pi^N : N \rightarrow \Sigma$  is the composition of the projection in the universal cover to the horizontal factor and then  $\pi^M$ .

When  $X$  is identically zero the submanifold  $M$  is horizontal in  $N$ , isometric to  $\Sigma$  by (3), the holonomy of the normal bundle  $\nu\Sigma$  is trivial, and, again,  $V$  is isometric to the direct product  $\Sigma \times (R^2, h)$  of the soul  $\Sigma$  and some non-negatively curved plane  $(R^2, h)$ .

Next we prove that if  $X$  is not identically zero, or has no zeros at all, then  $X$  vanish along some closed geodesic.

**Theorem 1.** *If the set of zeros  $\mathcal{Z}$  of the vector field  $X$  in  $M$  is a proper subset (i.e., is not  $M$  itself or empty) then  $\mathcal{Z}$  is a closed geodesic. Every minimal geodesic connecting two points from  $\mathcal{Z}$  is itself a subset of  $\mathcal{Z}$ .*

*Proof.* If  $\mathcal{Z}$  is a proper subset of  $M$  then for some  $P \in \mathcal{Z}$  there exists a sequence of points  $Q_i \rightarrow P$  such that  $X(Q_i) \neq 0$ . As in [GT], see Lemma 2.1; we note that every geodesic  $L(P, Q; t)$  in  $M$  connecting a point  $P$  where  $X$  vanish with an arbitrary point  $Q$  with non-vanishing  $X(Q)$  is orthogonal to  $X(Q)$ ,

$$(4) \quad \bar{Q}P \perp X(Q),$$

where  $\bar{P}Q$  denotes the vector of direction of  $L(P, Q; t)$  at the point  $Q$ . Hence,  $\mathcal{Z}$  belongs to the exponential image  $\Pi(Q, X(Q))$  of a plane in  $T_Q M$  of all vectors normal to  $X(Q)$ :

$$(5) \quad \mathcal{Z} \subset \Pi(Q, X(Q)).$$

The surface  $\Pi(Q, X(Q))$  near  $Q$  is "almost a plane" - smooth with a second form vanishing at  $Q$ . Fix for a moment some  $Q = Q_i$  close enough to  $P$ . Then in a small closed ball  $B$  around  $P$  with radius  $\text{dist}(P, Q)$  zeros of  $X$  belong to this "almost a plane"  $\Pi(Q, X(Q))$ . Thus there exists the farthest point  $Q' \in B$  to  $\Pi$  where  $X(Q') \neq 0$ . Then by (5) the part of the set  $\mathcal{Z}$  inside the ball  $B$  belongs to the intersection of two "almost planes"  $\Pi(Q, X(Q))$  and  $\Pi(Q', X(Q'))$  which are "almost orthogonal". This intersection, as easy to see, is a smooth curve with geodesic curvature of the order  $\text{dist}(P, Q)$ . Because the point  $Q = Q_i$  can be chosen arbitrary close to  $P$  we conclude that the set  $\mathcal{Z}$  is inside some finite collection of intervals of geodesics. Next, we verify that  $\mathcal{Z}$  is connected. Indeed, if not we may find two different points  $P_1$  and  $P_2$  from its different components and such that the minimal geodesic  $L = L(P_1, P_2; t)$  connecting these points does not intersect  $\mathcal{Z}$ . In some small ball  $B$  around the middle point  $Q$  of this geodesic the vector field  $X$  will be non-zero with  $X(Q)$  normal to  $L$ . Consider the "almost plane"  $\Pi$  in  $B$  going through the point  $Q$  and normal to  $L$ . From (4) we see that the vector field  $X$  in this plane not only is almost tangent to  $\Pi$ , but also almost tangent to small circles in  $\Pi$  around  $Q$ . Which implies that the projection of  $X$  on  $\Pi$  has index  $\pm 1$  at the center  $Q$  of these circles, i.e., equals zero at  $Q$ . The obtained contradiction proves that  $\mathcal{Z}$  is a closed geodesic.<sup>6</sup> Clearly, if some minimal geodesic  $L$  connects two zeros  $P$  and  $Q$  from  $\mathcal{Z}$ , but does not belong to  $\mathcal{Z}$  we may repeat arguments above to show that there exists one more point in the interior of  $L$  where  $X$  vanish. Which completes the proof of our theorem.

Note, that in our arguments we used only condition (4). Hence we have the following.

---

<sup>6</sup>By the arguments above we immediately deduce that  $\mathcal{Z}$  is a connected geodesic. The fact that this geodesic can not be infinite follows from the compactness of  $M$  and that it can not accumulate to something other than itself.

**Corollary 1.** *If some smooth (not identically zero) vector field  $X$  in some compact three-dimensional manifold  $M$  satisfies (4), then its set of zeros  $\mathcal{Z}$  is a closed geodesic.*

Next we consider the metric structures of  $M$  when  $\mathcal{Z}$  is a closed geodesic.<sup>7</sup>

### 3. $\mathcal{Z} = S^1$

Assume that the set of zeros  $\mathcal{Z}$  of the vector field  $X$  is some closed geodesic  $\mathcal{Z} = \mathcal{Z}(t), 0 \leq t \leq 1$ . Then  $\mathcal{Z}(t)$  is horizontal, its projection by the submersion  $\pi$  to the soul  $\Sigma$  is again a closed geodesic  $\bar{\mathcal{Z}}(t)$  of the same length, for every  $t$  the geodesic  $l_t(s)$  connecting  $\bar{\mathcal{Z}}(t)$  and  $\mathcal{Z}(t)$  is normal to  $\Sigma$  with a direction  $V(t)$  parallel along  $\bar{\mathcal{Z}}(t)$ . Now take an arbitrary point  $Q$  in  $M$  and connect it with all the points  $\mathcal{Z}(t)$  by minimal geodesics  $L_t(s)$ . All this geodesics are horizontal, and, if  $\bar{Q} = \pi(Q)$ , their projections  $\bar{L}_t(s)$  are minimal geodesics connecting  $\bar{Q}$  with  $\bar{\mathcal{Z}}(t)$ . Also, if  $V(t, s)$  denotes the (unit) vector of the direction of the (vertical) geodesic connecting  $\bar{L}_t(s)$  with  $L_t(s)$ , then  $V(t, s)$  is parallel along  $\bar{L}_t(s)$ . In particular, it follows that the parallel translation along a closed path from  $\bar{Q}$  to  $\bar{Q}$  consisting of two  $\bar{L}_{t'}(s)$  and  $\bar{L}_{t''}(s)$  and a part  $\bar{L}(t), t' \leq t \leq t''$  acts trivially on  $V$  - the direction of  $\bar{Q}Q$ . Which by the prism construction from [M1-3] implies that the O'Neill's fundamental tensor  $A$  vanishes at  $Q$  for horizontal vectors tangent to the family of geodesics  $L_t(s)$ . As we already saw, this family belongs to the plane  $\Pi(Q, X(Q))$  of all geodesics, issuing from  $Q$  in directions normal to  $X(Q)$ . Thus we have

$$(6) \quad A_Y Z(Q) \equiv 0$$

for all  $Y, Z$ . Again, by the same prism construction we have

$$(7) \quad R(\bar{Y}(t), \bar{Z}(t))V(t) \equiv 0$$

along  $\bar{\mathcal{Z}}(t)$ , where  $\bar{Z}(t)$  is the unit tangent to  $\bar{\mathcal{Z}}(t)$ ,  $\bar{Y}(t)$  any tangent to  $\Sigma$ , and  $V(t)$  is the direction of the vertical geodesic  $\bar{\mathcal{Z}}(t)\mathcal{Z}(t)$ .

Because  $Q$  (and  $\bar{Q}$  correspondingly) was arbitrary, the tensor  $A$  vanishes identically in  $M$  on vectors normal to the vector field  $X$ .

**Theorem 2.** *Distribution in  $M \setminus \mathcal{Z}$  of the two-planes normal to the vector field  $X$  is integrable. It is tangent to the family of totally geodesic spheres with a common intersection set - the closed geodesic  $\mathcal{Z}$ .*

*Proof.* Indeed, from (6) immediately follows that the Lie bracket of arbitrary fields  $Y, Z$  orthogonal to  $X$  is also orthogonal to  $X$ :

$$(8) \quad ([Y, Z], X) = ([Y, Z], W - (W, E)E) = ([Y, Z], W) = (A_Y Z - A_Z Y) = 0.$$

---

<sup>7</sup>When  $X$  has no zeros we may introduce the following **A-"contact" structure** on  $M$ . Denote by  $\alpha$  the 1-form on  $M^3$  given by the scalar product with a vector field  $X$ . Because the vector field  $E$  on  $N$  is parallel, from  $\nabla^N E \equiv 0$  (here  $\nabla^N$  is the covariant derivative in  $N$  in a metric induced by  $N \subset V$ ) we see that

$$(\nabla_Y^N X, Z) = (\nabla_Y W - (\nabla_Y W), E)E, Z) = (\nabla_Y W, Z)$$

for arbitrary  $Y, Z$  tangent to  $M$ . Therefore, because  $M$  is totally geodesic in  $N$  we have by direct calculations that

$$\alpha \wedge d\alpha = a(P) d\text{vol}^M,$$

where  $d\text{vol}^M$  is the volume form of  $M$  and the function  $a(P)$  is given by

$$a(P) = (A_Y Z, X)$$

or by  $(A'_Y Z', X')$  where  $\{X', Y', Z'\}$  an arbitrary orthonormal (positively orientated) basis in  $T_P M$ . The horizontal distribution on  $M$  is not involutive outside zeros of  $a$ . It would be interesting to find examples with  $a$  vanishing somewhere and nowhere zero  $\alpha$ .

Thus, the vector field  $X$  is, actually, the field of normals to some family of hyper-surfaces in  $M$ . But, as we already know, every geodesic  $L_t$  connecting  $Q$  with a point  $\mathcal{Z}(t)$  of  $\mathcal{Z}$  belongs to such a surface. From which, obviously, follows that this family of surfaces coincide with the family of our planes  $\Pi(Q, X(Q))$ . Because at  $Q$  the second form of this surface vanish, and  $Q$  is arbitrary our surfaces have vanishing second forms or are totally geodesic. For definiteness, from now on we call by  $\Pi(Q)$  the union of all geodesics  $L_t$  connecting  $Q$  with  $\mathcal{Z}$ . It is a totally geodesic surface which boundary is the closed geodesic  $\mathcal{Z}(t)$ . Therefore, the vector  $Y(t)$  tangent to  $\Pi$  and normal to this boundary at the point  $\mathcal{Z}(t)$  is parallel along  $\mathcal{Z}$ . Because the tangent vector  $Z(t)$  to this geodesic is also (auto-)parallel, we see that the holonomy around  $\mathcal{Z}$  is trivial, i.e., parallel translation along  $\mathcal{Z}(t)$  is the identity operator. If we choose some parallel vector field  $Y^*(y)$  along  $\mathcal{Z}$  normal to  $Z(t)$ , we can define the angle function  $\psi$  for vectors  $Y(t)$  normal to  $Z(t)$  as the angle between  $Y(t)$  and  $Y^*(t)$ . Corresponding  $\Pi(Q)$  we denote also by  $\Pi(\psi)$ . To complete the proof of the theorem we note that for (a half-sphere)  $\Pi(\psi)$  there exists another one  $\Pi(\psi + \pi)$  which normal to their common boundary  $\mathcal{Z}(t)$  equals  $-Y(t)$ . Their union in a neighborhood of  $\mathcal{Z}(t)$  is again an exponential image of planes tangent to  $Y(t)$  and  $Z(t)$ , and therefore, is a smooth surface: a sphere which we denote by  $S^2(Q)$ , or by  $S^2(\psi)$  (then  $S^2(\psi) = S^2(\psi + \pi)$ ). Theorem 2 is proved.

Configuration we described in the last theorem is well-known and is called **an open book decomposition**.

**Corollary 2.** *If the set of zeros of  $X$  is a closed geodesic  $\mathcal{Z}(t)$ , then  $M$  admits an open book decomposition with a bending  $\mathcal{Z}$  and pages  $\Pi(\psi)$  which are totally geodesic half-spheres.*

Next we look more closely on the family of diffeomorphisms between pages  $\Pi(\psi)$  of our open book decomposition given by shifts in directions normal to them. Let  $f_\theta : \Pi(\psi) \rightarrow \Pi(\psi + \theta)$  denotes the map sending the point  $Q$  in  $\Pi(\psi)$  into the intersection of  $\Pi(\psi + \theta)$  with an integral curve of the field of normals to pages issuing from  $Q$ . If we denote

$$(9) \quad \partial f_\theta(Q) / \partial \theta = X^*(f_\theta(Q)),$$

then the field  $X^*$  is proportional to  $X$ , i.e.,  $X^*(Q) = k'(Q)X(Q)$  for some positive function on  $M \setminus \mathcal{Z}$ . By  $k$  we denote its norm:  $k(Q) = \|X^*(Q)\|$ . Because all pages  $\Pi(\psi)$  are totally geodesic all maps  $f_\theta$  are isometries. Therefore, we call the family of these isometries: "turning pages". If, in addition,  $k$  is constant along trajectories of  $X^*$  (or  $X$ , which is the same) then  $f_\theta : M \rightarrow M$  is a family of isometries of the entire  $M$ ,<sup>8</sup> and the vector field  $X^*$  is a Killing vector field. Note also the following trivial statement.

**Lemma 3.** *All trajectories of the vector field  $X^*$  in  $M$  are closed circles around  $\mathcal{Z}$ .*

*Proof.* Indeed, take some geodesic  $L_t(s)$  in  $S^2(\psi)$  connecting some  $Q$  with the point  $\mathcal{Z}(t)$  which is nearest to it, and consider the orbit of this geodesic under our family of "rotations":  $\Phi(s, \theta) = f_\theta(L_t(s))$ . We choose natural parameter  $s$  on  $L_t(s)$  in such a way that  $\mathcal{Z}(t) = L_t(0)$ . Because for every  $\theta$  the curve  $f_\theta(L_t(s))$  lies in the totally geodesic  $\Pi(\psi + \theta)$  and  $f_\theta$  is an isometry, this  $f_\theta(L_t(s))$  is again the geodesic in  $M$ . Therefore,  $\Phi(s, \theta)$  is a part of the "plane"  $\Pi(\mathcal{Z}(t), Z(t))$  of all geodesics issuing from  $\mathcal{Z}(t)$  in directions normal to  $Z(t)$ . For a fixed  $s$  the line  $\Phi(s, \theta)$  is a closed circle in this "plane".<sup>9</sup>

---

<sup>8</sup>"reading the book"

<sup>9</sup>It is interesting to note also the following property of these circles. As we will show, the vector field  $X^*$  is Killing and constant along its trajectories, i.e., circles  $f_\theta(Q)$ . Therefore, the norm  $k$  of  $X^*$  attains its maximum on some set  $\mathcal{Z}^*$  which is invariant under rotations  $f_\theta$ . We claim that  $\mathcal{Z}^*$  is a collection of closed geodesics in  $M$ . Indeed, from  $Y(X^*, X^*) \equiv 0$  for every  $Y$  tangent to  $S^2(\psi)$  at some point  $Q$  of  $\mathcal{Z}^*$  it follows that  $\nabla_{X^*} X^* \equiv 0$ , or that the geodesic curvature of the orbit  $f_\theta(Q)$  equals zero. Every closed geodesic from  $\mathcal{Z}^*$  is linked with  $\mathcal{Z}$ .

#### 4. $X^*$ is Killing

As we saw above,  $A_Z$  vanishes along  $\mathcal{Z}$ , see (6,7). Also the holonomy of the normal bundle is trivial along the projection  $\bar{\mathcal{Z}}(t)$  of  $\mathcal{Z}$  under submersion  $\pi$  which is the closed geodesic in the soul  $\Sigma$ . Therefore, applying the simplified version of arguments<sup>10</sup> from the proof of the Theorem A from [M5] (see section 5 there) we get

$$(10) \quad \nabla_W W \equiv 0 \quad \text{and} \quad R[W(t), Z(t)] \equiv 0$$

for a unit vertical field  $W(t)$  along  $\mathcal{Z}$ .

Take another vector  $V(\phi)$  normal to  $\Sigma$  at  $\bar{P} = \bar{\mathcal{Z}}(0)$  with an angle  $\phi$  to  $V$ . Its parallel transport along  $\bar{\mathcal{Z}}(t)$  is again  $V(\phi)$ . Denote by  $V(\phi, t)$  the corresponding parallel vector field along  $\bar{\mathcal{Z}}(t)$ . The vertical lifts of  $\bar{\mathcal{Z}}(t)$  into  $N$  along this vector field are again closed geodesics which we denote by  $\mathcal{Z}(\phi, t)$ . Easy to see that (6,7) are satisfied along them<sup>11</sup>, which in turn implies (10) along  $\mathcal{Z}(\phi, t)$ , or that the vertical fibers of the submersion  $\pi : \mathcal{Z}(\phi, t) \rightarrow \bar{\mathcal{Z}}(t)$  have zero geodesic curvature, or are geodesic lines in  $N$ . Hence, they coincide with projections of straight lines, i.e.,  $R$ -factors under universal cover  $M \times R \rightarrow N$ . We formulate the obtained result as follows.

**Lemma 4.** *The set of zeros of the vector field  $X$  in  $N$  is a tori which is the image of the direct product of  $\mathcal{Z} \subset M$  with a straight-line factor  $R$  in the universal cover  $\tilde{N} = M \times R$  under covering map  $\tilde{N} \rightarrow N$ . For an arbitrary choice of  $M$  in  $N$  the  $\pi$ -projection of the set of zeros of  $X$  in  $M$  is the same closed geodesic  $\bar{\mathcal{Z}}$  in  $\Sigma$ .*

The obtained claim means that every vertical fibre in  $N$  stays in the set of zeros of the vector field  $X$  if it contains some of the point where  $X$  vanish. Now we can repeat arguments from [GT] and prove the following statement.

**Theorem 3.** *The vector field  $X^*$  is Killing, if it has non-empty zero set.*

*Proof.* Indeed, the Lemma 4's claim enable us to repeat arguments from [GT]: for every point  $Q$  of  $M$  denote by  $(f_t(Q), t)$  the points of the fiber of the submersion  $\pi : N \rightarrow \Sigma$  issuing from  $Q$ . These are trajectories of the vector field  $W$  in  $N$ . As we saw, the distance between  $f_t(Q)$  and  $f_t(P)$  is constant for every  $P$  from the zero set  $\mathcal{Z}$  since the geodesic connecting them in  $M \times \{t\}$  is horizontal. By the Lemma 4 we see  $f_t$  is the identity map on  $\mathcal{Z}$ . Therefore,  $f_t(Q)$  are circles  $S_Q^1$  around  $\mathcal{Z}$ , they coincide with the circles  $f_\theta(Q)$  which are orbits of the vector field  $X^*$  above. To show that  $X^*$  is Killing consider the cylinder  $C_Q = \{(f_\theta(Q), t)\}$ , (see Lemma 2.1 in [GT]). The restriction of  $\pi$  on  $C_Q$  is a Riemannian submersion of a flat cylinder onto some circle in  $\Sigma$ , or by [GG] has fibers tangent to some Killing field on  $C_Q$ . This proves that  $X^*$  has constant norm along  $C_Q$  and is a Killing vector field.

#### 5. Proof of the Theorem B

From Theorem 3 it follows that the restriction of the Riemannian submersion  $\pi : V \rightarrow \Sigma$  on  $N$ , which is the boundary of some  $s$ -metric neighborhood of the soul, can be described as the factoring by the action along trajectories of the Killing vector field  $X^*$ . From this fact the Theorem B follows in the same way as Theorem A; see section 3 in [GT] for the meticulous analysis of the cooperation between Killing vector fields  $X^*$  on different  $s$ -metric neighborhoods of the soul which ensures the claim of both Theorems A and B.

#### REFERENCES

- [CaS] J. Cao, M.-C. Shaw The smoothness of Riemannian submersions with nonnegative sectional curvature, arXiv: math.DG/0309328 (2003).

<sup>10</sup>when  $\text{codim}\Sigma = 2$  instead of  $\text{codim}\Sigma = 3$ .

<sup>11</sup>for the proof note, that (7) implies (6) through the prism construction, see the Lemma 2 in [M5]

- [CG] J. Cheeger, D. Gromoll, *On the structure of complete manifolds of nonnegative curvature*, Ann. Math. **96 no.3** (1972), 413–443.
- [GG] D. Gromoll and K. Grove, *One-dimensional metric foliations in constant curvature spaces*, Differential Geometry and Complex Analysis (1985), 165–168.
- [GT] D. Gromoll and K. Tapp, *Nonnegatively curved metrics on  $S^2 \times R^3$* , Geometriae Dedicata **99** (2003), 127–136.
- [M1] V. Marenich, *Metric structure of open manifolds of nonnegative curvature*, Doklady Acad. Sc. USSR **261:4** (1981), 801–804.
- [M1 rus] V. Marenich, *Metric structure of open manifolds of nonnegative curvature (complete version in russian)*, Ukrainian Geom. Sb. **26** (1983), 79–96.
- [M2] V. Marenich, *The metric of nonnegative curvature on the tangent bundle of two-dimensional sphere*, Sibirsk. Math. Zh. **27:2** (1986), 121–138.
- [M3] V. Marenich, *The holonomy in open manifolds of nonnegative curvature*, MSRI, Preprint No. 003–94. (1993).
- [M4] V. Marenich, *The holonomy in open manifolds of nonnegative curvature*, Michigan Math. Journal **43:2** (1996), 263–272.
- [M5] V. Marenich, *Rigidity of non-negatively curved metrics on open five-dimensional manifolds*, arXiv math/DG 0411632 (2004).
- [O’N1] B. O’Neill, *The fundamental equations of submersion*, Mich. Math. J. **13 no. 4** (1966), 459–469.
- [P] G. Perelman, *Proof of the soul conjecture of Cheeger and Gromoll*, J. Differential Geometry **40** (1994), 209–212.
- [T] K. Tapp, *Rigidity for Nonnegatively curved Curved Metrics on  $S^2 \times R^3$* , Ann. of Global Analysis and Geometry **25** (2004), 43–58.

HÖGSKOLAN I KALMAR, 391 82, KALMAR, SWEDEN

E-mail address: valery.marenich@hik.se, mikael.bengtsson@hik.se